



# First report of the *East African cassava mosaic virus-Uganda (EACMV-UG)* infecting cassava (*Manihot esculenta*) in Cameroon

S.A. Akinbade<sup>1</sup>, R. Hanna<sup>2</sup>, A. Nguenkam<sup>2</sup>, E. Njukwe<sup>2</sup>, A. Fotso<sup>2</sup>, A. Doumtsop<sup>2</sup>, J. Ngeve<sup>3</sup>, S.T.N. Tenku<sup>3</sup> and P. Lava Kumar<sup>1\*</sup>

<sup>1</sup> International Institute of Tropical Agriculture (IITA), PMB 5320, Ibadan, Nigeria ; <sup>2</sup> IITA, BP 2008 (Messa), Yaoundé, Cameroon ; <sup>3</sup> Institut de Recherche Agricole pour le Développement, BP 2123, Yaoundé, Cameroon

\*E-mail: l.kumar@cgiar.org

Received: 18 Jan 2010. Published: 12 May 2010.

Cassava mosaic disease (CMD) is a major constraint of cassava (*Manihot esculenta*) in Cameroon. Previous studies confirmed the occurrence of *African cassava mosaic virus* (ACMV), *East African cassava mosaic Cameroon virus* (EACMCV) and *East African cassava mosaic virus* (EACMV) in CMD etiology (Fondong *et al.*, 2000). In this study, surveys were conducted for EACMV-Uganda (EACMV-UG), a recombinant virus responsible for severe CMD epidemics in East and Central Africa, which occurs in adjoining Republic of Congo (RC) and Gabon (Legg *et al.*, 2004). Surveys were conducted during March to November 2009 in 156 fields in 37 administrative divisions (Fig. 1). Six to eight samples from each field were collected from plants with distinct symptoms and severity, and analysed with the polymerase chain reaction (PCR) using primers UV-AL1/F1 and ACMV-CP/R3 specific for EACMV-UG (Zhou *et al.* 1997), ACMV and EACMV/ EACMCV (Alabi *et al.*, 2008). Viruses were identified based on the amplification of diagnostic fragments for ACMV (368 bp), EACMV/EACMCV (650 bp), and EACMV-UG (1630 bp).

Out of 1159 samples, single infections of ACMV and EACMV/EACMCV, and mixed infections of ACMV with EACMV/EACMCV were detected in, respectively, 57.1%, 4.2% and 23.0% of the total. EACMV-UG, as a mixed infection with ACMV (Fig. 2) and/or EACMV/EACMCV, was detected in 57 (33.9%) out of 168 samples from a total of 21 fields in Kadey, Boumba-and-Ngoko, Lom-and-Djerem and Mbere divisions (Figs. 1). Incidence in fields ranged from 11.1% (Mbotoro) to 83.3% (Boyere). To further confirm the virus identity, the 1630 bp PCR product from Gamago (Kadey) was sequenced (GenBank Accession No. GU395301). Comparison with sequences in GenBank revealed the highest identity (98%) with the DNA-A sequence of EACMV-UG (Fig. 3). To our knowledge, this is the first report of EACMV-UG in Cameroon. Its occurrence in areas bordering EACMV-UG affected countries suggests the

spread of the virus by whiteflies or infected cuttings. Urgent deployment of resistant varieties in the affected regions is necessary to prevent severe CMD epidemics and further spread of this virus into West Africa.

## Acknowledgements

This research was supported with funds provided to IITA by the International Fund for Agricultural Development (IFAD) and by IITA core donors.

## References

- Alabi OJ, Kumar PL, Naidu RA, 2008. Multiplex PCR method for the detection of *African cassava mosaic virus* and *East African cassava mosaic Cameroon virus* in cassava. *Journal of Virological Methods* **154**, 111-120. [doi:10.1016/j.jviromet.2008.08.008]
- Fondong VN, Pitta, JS, Rey MEC, de Kochko A, Beachy RN, Fauquet CM, 2000. Evidence of synergism between African cassava mosaic virus and a new double-recombinant geminivirus infecting cassava in Cameroon. *Journal of General Virology* **81**, 287-297.
- Legg JP, Ndjelassili F, Okao-Okuja G, 2004. First report of cassava mosaic disease and cassava mosaic geminiviruses in Gabon. *Plant Pathology* **53**, 232. [doi:doi:10.1111/j.0032-0862.2004.00972.x]
- Zhou X, Liu Y, Calvert L, Munoz C, Otim-Nape GW, Robinson DJ, Harrison BD, 1997. Evidence that DNA A of a geminivirus associated with severe cassava mosaic disease in Uganda has arisen by interspecific recombination. *Journal of General Virology* **78**, 2101-2111.




Figure 1



Figure 2




Figure 3

**To cite this report:** Akinbade SA, Hanna R, Nguenkam A, Njukwe E, Fotso A, Doumtsop A, Ngeve J, Tenku STN, Lava Kumar P, 2010. First report of the *East African cassava mosaic virus-Uganda* (EACMV-UG) infecting cassava (*Manihot esculenta*) in Cameroon. *New Disease Reports* **21**, 22. [doi:10.5197/j.2044-0588.2010.021.022]

©2010 The Authors

This report was published on-line at [www.ndrs.org.uk](http://www.ndrs.org.uk) where high quality versions of the figures can be found.